

- 24) Anthes E. Evidence-based medicine: Save blood, save lives. *Nature* 2015; **520**: 24-6.
- 25) Gouezec H, Berger E, Bergoin-Costello V, et al; Groupe des Hémobiologistes et Correspondants d'Hémovigilance. [Suitability of red blood cell transfusion: a multicenter study.] *Transfus Clin Biol* 2010; **17**: 318-30. [In French.]
- 26) Serious Hazards of Transfusion Steering Committee. Serious hazards of transfusion: annual report 2014. Available at: <http://www.shotuk.org/shot-reports/report-summary-supplement-2014/>. Accessed on 10/12/2015.
- 27) Villanueva C, Colomo A, Bosch A, et al. Transfusion strategies for acute upper gastrointestinal bleeding. *N Engl J Med* 2013; **368**: 11-21.
- 28) Leal-Noval SR, Muñoz-Gómez M, Jiménez-Sánchez M, et al. Red blood cell transfusion in non-bleeding critically ill patients with moderate anemia: is there a benefit? *Intensive Care Med* 2013; **39**: 45-53.
- 29) Ferraris VA, Davenport DL, Saha SP, et al. Surgical outcomes and transfusion of minimal amounts of blood in the operating room. *Arch Surg* 2012; **147**: 49-55.
- 30) Grey DE, Finlayson J. Red cell transfusion for iron deficiency anaemia: a retrospective audit at a tertiary hospital. *Vox Sang* 2008; **94**: 138-42.
- 31) Joshi G, Carroll M, O'Rourke P, Coffey F. Role of quality assessment in improving red blood cell transfusion practice. *Ir J Med Sci* 1997; **166**: 16-9.
- 32) Gillham M, Mark A. A retrospective audit of blood loss in total hip joint replacement surgery at Middlemore Hospital. *N Z Med J* 1997; **110**: 294-7.
- 33) Allameddine A, Heaton M, Jenkins H, et al. Inappropriate use of blood transfusion in emergency department in a tertiary care hospital and potential saving with patient blood management [abstract]. *Transfus Med* 2014; **24**: 25.
- 34) Quintana-Díaz M, Fabra-Cadenas S, Gómez-Ramírez S, et al. A fast-track anaemia clinic in the Emergency Department: feasibility and efficacy of intravenous iron administration for treating sub-acute iron deficiency anaemia. *Blood Transfus* 2016; **14**: 126-33.
- 35) Tuckfield A, Haeusler MN, Rigg AP, Metz J. Reduction of inappropriate use of blood products by prospective monitoring of transfusion request forms. *Med J Aust* 1997; **167**: 73-6.
- 36) Jackson GN, Snowden CA, Indrikovs AJ. A prospective audit program to determine blood component transfusion appropriateness at a large university hospital: a 5-year experience. *Transfus Med Rev* 2008; **22**: 154-61.
- 37) Goodnough LT. Trends in blood utilization. *Transfus Med* 2014; **24**: 2.
- 38) Rohde JM, Dimcheff DE, Blumberg, et al. Health care-associated infection after red blood cell transfusion: a systematic review and meta-analysis. *JAMA* 2014; **311**: 1317-26.
- 39) Audet AM, Goodnough LT, Parvin CA. Evaluating the appropriateness of red blood cell transfusions: the limitations of retrospective medical record reviews. *Int J Qual Health Care* 1996; **8**: 41-9.
- 40) Marvulo NL, Bonatto RC, Carpi MF, et al. [Red blood cell transfusion in children admitted in a pediatric intensive care unit.] *RBTI* 2006; **18**: 390-5. [In Portuguese.]
- 41) de Souza DA, Silva FG, Costa PJ. Critical evaluation of justifications for the transfusion of red blood cells: the reality of a government emergency hospital. *Rev Bras Hematol Hemoter* 2013; **35**: 263-7.
- 42) Friedman MT, Ebrahim A. Adequacy of physician documentation of red blood cell transfusion and correlation with assessment of transfusion appropriateness. *Arch Pathol Lab Med* 2006; **130**: 474-9.

Arrived: 8 December 2015 - Revision accepted: 4 April 2016

Correspondence: Manuel Muñoz

Perioperative Transfusion Medicine - School of Medicine, University of Málaga
Campus de Teatinos
29071 Málaga, Spain
e-mail: mmunoz@uma.es

Appendix I

Researchers of the group for the Safe, Effective and Efficient Blood Use in Emergency Departments USEES-URG (in alphabetical order):

- Borobia, Alberto M (Department of Clinical Pharmacology, La Paz University Hospital);
- Carcas-Sansuan, Antonio José (Department of Clinical Pharmacology, La Paz University Hospital);
- Fabra, Sara (Emergency Department, La Paz University Hospital);
- Frías-Iniesta, Jesús (Department of Clinical Pharmacology, La Paz University Hospital);
- García-Erce, José A (Service of Haematology and Haemotherapy, San Jorge Hospital, Huesca, Aragon Institute of Health Science (IACS); investigator;
- Garde, Gregorio (Emergency Department, Virgen de la Luz Hospital);
- Guallar, Camino (Intensive Care Unit, La Paz University Hospital);
- Gutiérrez, Gabriel (Emergency Department, Hospital of Cruces);
- Lozano, Consuelo (Emergency Department, Nuestra Señora del Prado Hospital);
- Marcos, Faustino A (Emergency Department, Nuestra Señora del Prado Hospital);
- Maroun-Eid, Charbel (Emergency Department, La Paz University Hospital);
- Medrano, Nicolás (Department of Clinical Pharmacology, La Paz University Hospital);
- Millán, Pablo (Intensive Care Unit, La Paz University Hospital);
- Muñoz, Manuel (Department of Surgical Specialties, Biochemistry and Immunology, School of Medicine, University of Málaga, Málaga);
- Panadero Sánchez, Alicia (Emergency Department, Virgen de la Luz Hospital);
- Pozuelo, Itziar (Intensive Care Unit, La Paz University Hospital);
- Quintana-Díaz, Manuel (Emergency Department, La Paz University Hospital);
- Ramírez Elena (Department of Clinical Pharmacology, La Paz University Hospital);
- Reche, Begoña (Emergency Department, La Paz University Hospital);
- Rodríguez de la Rúa, Ana (Department of Haematology, La Paz University Hospital);
- Rodríguez Villa, Sancho (Emergency Department, Hospital of Toledo);
- Sánchez-Casado, Marcelino (Emergency Department, Virgen de la Salud Hospital);
- Tong, Hoi Y (Department of Clinical Pharmacology, La Paz University Hospital);
- Viejo, Aurora (Department of Haematology, La Paz University Hospital).